



ORF 307: Lecture 6

Linear Programming: Chapter 5 Duality

Robert Vanderbei

Feb 21, 2019

Slides last edited on February 21, 2019

**ORF ENTERTAINMENT
THURSDAY
FEBRUARY 28TH
7:30 PM
SHERRERD HALL ATRIUM**

ALL ARE WELCOME, PRIZES, FOOD, FUN

**PERFORMANCE TIMES ARE STILL
AVAILABLE.... IT'S NOT TOO LATE TO
SIGN UP!!!**

NEW THIS YEAR KARAOKE!!!

CONTACT TZGLER@PRINCETON.EDU



Duality

Every Problem:

$$\begin{array}{ll}\text{maximize} & \sum_{j=1}^n c_j x_j \\ \text{subject to} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad i = 1, 2, \dots, m \\ & x_j \geq 0 \quad j = 1, 2, \dots, n,\end{array}$$

Has a Dual:

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^m b_i y_i \\ \text{subject to} & \sum_{i=1}^m y_i a_{ij} \geq c_j \quad j = 1, 2, \dots, n \\ & y_i \geq 0 \quad i = 1, 2, \dots, m.\end{array}$$

Dual of Dual

Primal Problem:

$$\begin{array}{ll}\text{maximize} & \sum_{j=1}^n c_j x_j \\ \text{subject to} & \sum_{j=1}^n a_{ij} x_j \leq b_i \quad i = 1, \dots, m \\ & x_j \geq 0 \quad j = 1, \dots, n\end{array}$$

Original problem is called the *primal problem*.

A problem is defined by its data (notation used for the variables is arbitrary).

Dual in “Standard” Form:

$$\begin{array}{ll}\text{—maximize} & \sum_{i=1}^m -b_i y_i \\ \text{subject to} & \sum_{i=1}^m -a_{ij} y_i \leq -c_j \quad j = 1, \dots, n \\ & y_i \geq 0 \quad i = 1, \dots, m\end{array}$$

Dual is “negative transpose” of primal.

Theorem *Dual of dual is primal.*

Weak Duality Theorem

If (x_1, x_2, \dots, x_n) is feasible for the primal and (y_1, y_2, \dots, y_m) is feasible for the dual, then

$$\sum_j c_j x_j \leq \sum_i b_i y_i.$$


Proof.

$$\begin{aligned} \sum_j c_j x_j &\leq \sum_j \left(\sum_i y_i a_{ij} \right) x_j \\ &= \sum_{ij} y_i a_{ij} x_j \\ &= \sum_i \left(\sum_j a_{ij} x_j \right) y_i \\ &\leq \sum_i b_i y_i. \end{aligned}$$

Gap or No Gap?

An important question:

Is there a gap between the **largest primal value** and the **smallest dual value**?



Answer is provided by the Strong Duality Theorem (coming later).

Simplex Method and Duality

A Primal Problem:

$$\begin{aligned}
 \text{maximize } \zeta &= 0 + (-2)x_1 + 1x_2 + 1x_3 + 4x_4 \\
 w_1 &= 4 - 0x_1 - 1x_2 - 1x_3 - 2x_4 \\
 w_2 &= 3 - 0x_1 - 0x_2 - 2x_3 - 2x_4 \\
 w_3 &= 5 - 6x_1 - 2x_2 - (-2)x_3 - 5x_4
 \end{aligned}$$

Its Dual:

$$\begin{aligned}
 \text{maximize } -\xi &= 0 + (-4)y_1 + (-3)y_2 + (-5)y_3 \\
 z_1 &= 2 - 0y_1 - 0y_2 - (-6)y_3 \\
 z_2 &= -1 - (-1)y_1 - 0y_2 - (-2)y_3 \\
 z_3 &= -1 - (-1)y_1 - (-2)y_2 - 2y_3 \\
 z_4 &= -4 - (-2)y_1 - (-2)y_2 - (-5)y_3
 \end{aligned}$$

Notes:

- Dual is negative transpose of primal.
- Primal is feasible, dual is not.

Use primal to choose pivot: x_4 enters, w_3 leaves.

Make analogous pivot in dual: z_4 leaves, y_3 enters.

Second Iteration

After First Pivot:

$$\text{maximize } \zeta = 4 + \frac{-34}{5}x_1 + \frac{-3}{5}x_2 + \frac{13}{5}x_3 + \frac{-4}{5}w_3$$

Primal (feasible):

$$\begin{aligned} w_1 &= 2 - \frac{12}{5}x_1 - \frac{1}{5}x_2 - \frac{9}{5}x_3 - \frac{2}{5}w_3 \\ w_2 &= 1 - \frac{12}{5}x_1 - \frac{4}{5}x_2 - \frac{14}{5}x_3 - \frac{2}{5}w_3 \\ x_4 &= 1 - \frac{6}{5}x_1 - \frac{2}{5}x_2 - \frac{2}{5}x_3 - \frac{1}{5}w_3 \end{aligned}$$

$$\text{maximize } -\xi = -4 + -2y_1 + -1y_2 + -1z_4$$

Dual (still not feasible):

$$\begin{aligned} z_1 &= \frac{34}{5} - \frac{12}{5}y_1 - \frac{12}{5}y_2 - \frac{6}{5}z_4 \\ z_2 &= \frac{3}{5} - \frac{1}{5}y_1 - \frac{4}{5}y_2 - \frac{2}{5}z_4 \\ z_3 &= \frac{-13}{5} - \frac{9}{5}y_1 - \frac{14}{5}y_2 - \frac{2}{5}z_4 \\ y_3 &= \frac{4}{5} - \frac{2}{5}y_1 - \frac{2}{5}y_2 - \frac{1}{5}z_4 \end{aligned}$$

Note: *negative transpose property intact.*

Again, use primal to pick pivot: x_3 enters, w_2 leaves.

Make analogous pivot in dual: z_3 leaves, y_2 enters.

Third Iteration

After Second Pivot:

$$\text{maximize } \zeta = \boxed{69/14} + \boxed{-32/7} x_1 + \boxed{1/7} x_2 + \boxed{-13/14} w_2 + \boxed{-3/7} w_3$$

Primal (feasible):

$$\begin{aligned} w_1 &= \boxed{19/14} - \boxed{-6/7} x_1 - \boxed{5/7} x_2 - \boxed{-9/14} w_2 - \boxed{-1/7} w_3 \\ x_3 &= \boxed{5/14} - \boxed{-6/7} x_1 - \boxed{-2/7} x_2 - \boxed{5/14} w_2 - \boxed{-1/7} w_3 \\ x_4 &= \boxed{8/7} - \boxed{6/7} x_1 - \boxed{2/7} x_2 - \boxed{1/7} w_2 - \boxed{1/7} w_3 \end{aligned}$$

$$\text{maximize } -\xi = \boxed{-69/14} + \boxed{-19/14} y_1 + \boxed{-5/14} z_3 + \boxed{-8/7} z_4$$

Dual (still not feasible):

$$\begin{aligned} z_1 &= \boxed{32/7} - \boxed{6/7} y_1 - \boxed{6/7} z_3 - \boxed{-6/7} z_4 \\ z_2 &= \boxed{-1/7} - \boxed{-5/7} y_1 - \boxed{2/7} z_3 - \boxed{-2/7} z_4 \\ y_2 &= \boxed{13/14} - \boxed{9/14} y_1 - \boxed{-5/14} z_3 - \boxed{-1/7} z_4 \\ y_3 &= \boxed{3/7} - \boxed{1/7} y_1 - \boxed{1/7} z_3 - \boxed{-1/7} z_4 \end{aligned}$$

Note: *negative transpose property intact.*

Again, use primal to pick pivot: x_2 enters, w_1 leaves.

Make analogous pivot in dual: z_2 leaves, y_1 enters.

After Third Iteration

Primal:

$$\begin{aligned}
 \text{maximize } \zeta &= \boxed{26/5} + \boxed{-22/5} x_1 + \boxed{-1/5} w_1 + \boxed{-4/5} w_2 + \boxed{-2/5} w_3 \\
 x_2 &= \boxed{19/10} - \boxed{-6/5} x_1 - \boxed{7/5} w_1 - \boxed{-9/10} w_2 - \boxed{-1/5} w_3 \\
 x_3 &= \boxed{9/10} - \boxed{-6/5} x_1 - \boxed{2/5} w_1 - \boxed{1/10} w_2 - \boxed{-1/5} w_3 \\
 x_4 &= \boxed{3/5} - \boxed{6/5} x_1 - \boxed{-2/5} w_1 - \boxed{2/5} w_2 - \boxed{1/5} w_3
 \end{aligned}$$

- Is *optimal*.

Dual:

$$\begin{aligned}
 \text{maximize } -\xi &= \boxed{-26/5} + \boxed{-19/10} z_2 + \boxed{-9/10} z_3 + \boxed{-3/5} z_4 \\
 z_1 &= \boxed{22/5} - \boxed{6/5} z_2 - \boxed{6/5} z_3 - \boxed{-6/5} z_4 \\
 y_1 &= \boxed{1/5} - \boxed{-7/5} z_2 - \boxed{-2/5} z_3 - \boxed{2/5} z_4 \\
 y_2 &= \boxed{4/5} - \boxed{9/10} z_2 - \boxed{-1/10} z_3 - \boxed{-2/5} z_4 \\
 y_3 &= \boxed{2/5} - \boxed{1/5} z_2 - \boxed{1/5} z_3 - \boxed{-1/5} z_4
 \end{aligned}$$

- Negative transpose property remains intact.
- Is *optimal*.

Conclusion

Simplex method applied to primal problem (two phases, if necessary), solves both the primal and the dual.

Algebra of a Pivot

A primal pivot:

d	c	
b	a	

$\xrightarrow{\text{pivot}}$

$d - \frac{bc}{a}$	c/a	
$-b/a$	$1/a$	

The corresponding dual pivot:

$-d$	$-b$	
$-c$	$-a$	

$\xrightarrow{\text{pivot}}$

$-d + \frac{bc}{a}$	b/a	
$-c/a$	$-1/a$	

Strong Duality Theorem

Conclusion on previous slide is the essence of the *strong duality theorem* which we now state:

Theorem. *If the primal problem has an optimal solution,*

$$x^* = (x_1^*, x_2^*, \dots, x_n^*),$$

then the dual also has an optimal solution,

$$y^* = (y_1^*, y_2^*, \dots, y_m^*),$$

and

$$\sum_j c_j x_j^* = \sum_i b_i y_i^*.$$

Paraphrase:

If primal has an optimal solution, then there is no duality gap.

Duality Gap

Four possibilities:

- Primal optimal, dual optimal (no gap).
- Primal unbounded, dual infeasible (no gap).
- Primal infeasible, dual unbounded (no gap).
- Primal infeasible, dual infeasible (infinite gap).

\Leftarrow Strong Duality Theorem

\Leftarrow Weak Duality Theorem

\Leftarrow Weak Duality Theorem

See example below

Example of *infinite gap*:

$$\begin{array}{ll} \text{maximize} & 2x_1 - x_2 \\ \text{subject to} & x_1 - x_2 \leq 1 \\ & -x_1 + x_2 \leq -2 \\ & x_1, x_2 \geq 0. \end{array}$$